arXiv:math/0504317v2 [math.AP] 28 Jul 2005

Remarks on the Extremal Functions for the Moser-Trudinger Inequalities

Yuxiang Li
Department of Mathematical Science ,Tsinghua University, Beijing, P.R.China, 100084
E-mail address: yxli@math.tsinghua.edu.cn

Abstract

We will show in this paper that if λ is very close to 1, then

$$I(M,\lambda,m)=\sup_{u\in H_0^{1,n}(M),\int_M|\nabla u|^n dV=1}\int_\Omega(e^{\alpha_n|u|^{\frac{n}{n-1}}}-\lambda\sum_{k=1}^m\frac{|\alpha_n u^{\frac{n}{n-1}}|^k}{k!})dV,$$

can be attained, where M is a compact manifold with boundary. This result gives a counter example to the conjecture of de Figueiredo, do ó, and Ruf in their paper titled "On a inequality by N.Trudinger and J.Moser and related elliptic equations" (Comm. Pure. Appl. Math.,**55**:135-152, 2002).

Keywords: Moser-Trudinger inequality, extremal function.

MR (2000) Subject Classification: 58J05

1 Introduction:

Recall the Sobolev embedding: $H^{1,p}(\Omega)\hookrightarrow L^{\frac{np}{n-p}}(\Omega)$ where Ω is a bounded domain in $\mathbb{R}^n$ and $p\in(1,n)$.

When $p=n$, the critical growth is given by the Moser-Trudinger inequality ([5]) which can be expressed as:

$$\sup_{u\in H_0^{1,n}(\Omega),||\nabla u||_{L^n(\Omega)=1}}\int_\Omega(e^{\alpha|u|^{\frac{n}{n-1}}}-1)dx=c_n(\alpha)|\Omega|, \tag{1.2}$$

then, denoting by ω_{n-1} the measure of unit sphere in $\mathbb{R}^n$, one has

$$c_n(\alpha)<+\infty \ \ for \ \ 0<\alpha\leq\alpha_n=n\omega_{n-1}^{\frac{1}{n-1}},$$

$$c_n(\alpha)=+\infty \ \ for \ \ \alpha>\alpha_n.$$

Unlike Sobolev inequalities, the extremal functions for (1.2) with $\alpha=\alpha_n$ exist generally. The first result for the existence of extremal functions belongs to Carleson and Chang ([1]) who proved that (1.2) is attained when $\Omega=B_1(0)$ and $\alpha=\alpha_n$. Their work has been extended to bounded domains by Flucher([3]) and K-C, Lin([8]).

In [2] de Figueiredo et. al. gave a new proof for Carleson-Chang's theorem. In fact, they got a generalized result which states that

$$\sup_{u\in H_0^{1,n}(B_1(0)),||\nabla u||_{L^n(B_1(0))}=1}\int_{B_1(0)}(e^{\alpha_n|u|^{\frac{n}{n-1}}}-\lambda|u|^{\frac{n}{n-1}})dx$$

is attained for any $\lambda<\alpha_n$. Moreover, they proved the following:
Theorem A *let $F(t)=F(|t|)$ be a function increasing on $\mathbb{R}^+$ which satisfies*

$$1\leq F(t)\leq e^{\alpha_n|t|^{\frac{n}{n-1}}},\ and\ \lim_{t\to+\infty}\frac{F(t)}{e^{\alpha_n|t|^{\frac{n}{n-1}}}}=1,$$

then either

$$\sup_{u\in H_0^{1,n}(B_1(0)),\int_{B_1(0)}|\nabla u|^n=1}\int_{B_1(0)}F(u)dx$$

can be attained, or

$$\sup_{u\in H_0^{1,n}(B_1(0)),\int_{B_1(0)}|\nabla u|^n=1}\int_{B_1(0)}F(u)dx=e^{1+1/2+\cdots+1/(n-1)}+|B_1(0)|. \tag{1.3}$$

de Figueiredo et. al. then raised the following open problem in [1]:
Open problem: *Show that*

$$\sup_{\int_{B_1}|\nabla u|^n=1,u\in H_0^{1,2}(B_1)}\int_{B_1}F(u)dV$$

is not attained for $F(t)$ of the form

$$F(t)=e^{\alpha_n|t|^{\frac{n}{n-1}}}-g(t),$$

with

$$\lim_{t\to\infty}\frac{g(t)}{e^{\alpha_n|t|^{\frac{n}{n-1}}}}=0\ \ and\ \ g(t)\geq\alpha_n|t|^{\frac{n}{n-1}}.$$

However, this problem is much more complicated than they expected, we will show in this paper that the open problem is not true when $g(t)$ is very close to

$$g_m(t)=\sum_{k=1}^{m}\frac{|\alpha_n t^{\frac{n}{n-1}}|^k}{k!}.$$

Let

$$F_{\lambda,m}(t)=e^{\alpha_n|t|^{\frac{n}{n-1}}}-\lambda g_m(t),$$

and

$$I(\Omega,\lambda,m)=\sup_{u\in H_0^{1,n}(\Omega),\int_\Omega|\nabla u|dV=1}\int_\Omega F_{\lambda,m}(u)dV,$$

we shall prove that $I(\Omega,\lambda,m)$ is attained when λ is close to 1, in fact we shall prove it is attained even when Ω is a manifold.

The author have adopted blow up analysis to study a similar problem. In [7] the author extended Carleson-Chang's result to compact Riemannian manifolds. With the same method in [7], we can also get a generalization of Theorem A as follows:

Theorem 1.1: *Let (M,g) be a compact Riemannian manifold with boundary. If $I(M,\lambda,m)$ can not be attained, then we have*

$$I(M,\lambda,m)=\mu(M)+\frac{\omega_{n-1}}{n}e^{\alpha_n S_p+1+1/2+\cdots+1/(n-1)},$$

for some $p\overline{\in}\partial M$, where $\mu(M)$ is the measure of M.

Here we should explain what S_p is. Let G_p be the n-Lapalace Green function which is defined by

$$\begin{cases} -div(|\nabla G_p|^{n-2}\nabla G_p)=\delta_p \\ G_p|_{\partial M}=0. \end{cases} \tag{1.4}$$

By the asymptotic expansion theorem in [4] (the author has shown in [7] that their proof still works on manifolds), the function $G_p(x)+\frac{1}{\alpha_n}\log dist^n(x,p)$, is continuous at p. S_p is just its value at p, i.e.

$$S_p=\lim_{x\to p}(G_p(x)+\frac{1}{\alpha_n}\log dist^n(x,p)). \tag{1.5}$$

Since the proof for Theorem 1.1 is the same as the one done in [7] (also in [6]), we only give an outline of the proof from which one can see the idea why we construct the function sequence (3.5).

Then we will show that

Theorem 1.2: $I(M,1,m)>\mu(M)+\frac{\omega_{n-1}}{n}e^{\alpha_n S_p+1+1/2+\cdots+1/(n-1)}$.

Clearly, the function $f(\lambda)=I(M,\lambda,m)$ is continuous for fixed integer m, therefore we have

Main Theorem: *There is a constant $\lambda_0>1$, s.t. $I(M,\lambda,m)$ can be attained on $[0,\lambda_0)$.*

2 The outline of the proof for Theorem 1.1

Let $u_k\in H_0^{1,n}(M)$ which satisfies $u_k\geq 0$, $\int_M|\nabla u_k|^n dV_g=1$ and

$$\int_M(e^{\beta_k|u_k|^{\frac{n}{n-1}}}-\lambda g_m(u_k))dV_g=\sup_{v\in H_0^{1,n}(M),\int_M|\nabla v|^n dV_g=1}\int_M(e^{\beta_k|v|^{\frac{n}{n-1}}}-\lambda g_m(v))dV_g,$$

where $\{\beta_k\}$ is an increasing sequence which converges to α_n. Then we have

$$-div|\nabla u_k|^{n-2}\nabla u_k=\frac{u_k^{\frac{1}{n-1}}e^{\beta_k u_k^{\frac{n}{n-1}}}-\lambda g_m'(u_k)}{\lambda_k},$$

where

$$\lambda_k=\int_M(u_k^{\frac{n}{n-1}}e^{\beta_k u_k^{\frac{n}{n-1}}}-\lambda g_m' u_k)dV_g.$$

Let $c_k = \max\limits_M u_k(x)$, and $x_k \to p$. Then using the same method in [7], we have $p\overline{\in}\partial M$.

Take a normal coordinate system (U, x) around p. Let $r_k^n = \frac{\lambda_k}{c_k^{\frac{n}{n-1}} e^{\beta_k c_k^{\frac{n}{n-1}}}}$, then we have $r_k \to 0$ and

$$\frac{n}{n-1}\beta_k c_k^{\frac{1}{n-1}}(u_k(x_k + r_k x) - c_k) \to -n\log(1 + c_n r^{\frac{n}{n-1}}) \tag{2.1}$$

on any $B_L(0)$, where $c_n = (\frac{\omega_{n-1}}{n})^{\frac{1}{n-1}}$.

Moreover, we can get

$$c_k^{\frac{1}{n-1}} u_k \to G_p \tag{2.2}$$

on any $M \setminus B_\delta(p)$ where G_p is the Green function defined by (1.4).

It follows from (2.1) that

$$\sup_{v\in H_0^{1,2}(M), \int_M |\nabla v|^n dV_g = 1} F_{\lambda,m}(v) dV_g = \lim_{k\to+\infty}\int_M (e^{\beta_k |u_k|^{\frac{n}{n-1}}} - \lambda g_m(u_k)) dV_g = \mu(M) + \lim_{k\to+\infty}\frac{\lambda_k}{c_k^{\frac{n}{n-1}}}.$$

Then Theorem 1.1 can be obtained from an estimate for $\frac{\lambda_k}{c_k^{\frac{n}{n-1}}}$.

3 The proof of Theorem 1.2 :

The following lemma will play an important role in our computation.
Lemma 3.1 *Let $A_t = \{x \in M : G_p(x) > t\}$, then as $t \to +\infty$, we have*

$$\int_{\partial A_t} \frac{1}{|\nabla G_p|} \geq \omega_{n-1}^{\frac{n}{n-1}} e^{-\alpha_n t + \alpha_n S_p}(1 + O(e^{-\frac{2}{n}\alpha_n t})).$$

Proof: We have

$$-\frac{d|A_t|}{dt} = \int_{\partial A_t} \frac{dS_t}{|\nabla G_p|}. \tag{3.1}$$

By Hölder inequality, we get

$$(\int_{\partial A_t} \frac{dS_t}{|\nabla G_p|})^{n-1} \int_{\partial A_t} |\nabla G_p|^{n-1} dS_t \geq |\partial A_t|^n. \tag{3.2}$$

Since $\nabla G_p = \frac{\partial G_p}{\partial n}$ on ∂A_t, we have

$$\int_{\partial A_t} |\nabla G_p|^{n-1} dS_t = \int_{\partial A_t} |\nabla G_p|^{n-2} \frac{\partial G_p}{\partial n} \times 1 dS_t = 1.$$

Choose a normal coordinate system around p, then we are able to assume that $g = \sum\limits_i (dx^i)^2 + O(r^2)$, where $r^2 = dist^2(x, p) = \sum\limits_k (x^k)^2$.

Set

$$G_p(x) = -\frac{1}{\alpha_n}\log r^n + S_p + H(x),$$

where H is a continuous function with $H(0) = 0$. Then for a point $x \in \partial A_t$, we have $t = -\frac{1}{\alpha_n}\log r^n(x) + S_p + H(x)$, i.e.

$$r(x) = e^{-\frac{\alpha_n}{n}t + \frac{\alpha_n S_p + H(x)}{n}}. \tag{3.3}$$

We can easily see that

$$|A_t| = \int_{A_t} \sqrt{|g|}dx = \int_{A_t} (1 + O(r^2))dx = (1 + O(e^{-\frac{2\alpha_n}{n}t}))L(A_t),$$

where $L(\cdot)$ is the standard measure in $\mathbb{R}^n$. In the same way, we have

$$|\partial A_t| = (1 + O(e^{-\frac{2\alpha_n}{n}t}))L(\partial A_t).$$

Applying isopermetric inequality on $\mathbb{R}^n$, we get

$$|\partial A_t|^{\frac{n}{n-1}} \geq \alpha_n(1 + O(e^{-\frac{2\alpha_n}{n}t}))|A_t|.$$

Thus, by (3.1), (3.2) we have

$$-\frac{d|A_t|}{dt} \geq \alpha_n(1 + O(e^{-\frac{2}{n}\alpha_n t}))|A_t|. \tag{3.4}$$

Hence, we have

$$\frac{d(e^{\alpha_n t}|A_t|)}{dt} \leq O(e^{-\frac{2}{n}\alpha_n t})(e^{\alpha_n t}|A_t|) = O(e^{-\frac{2}{n}\alpha_n t}).$$

It is easy to see that $\lim\limits_{t\to\infty} e^{\alpha_n t}|A_t| = \frac{\omega_{n-1}}{n}e^{\alpha_n S_p}$, then we get

$$e^{\alpha_n t}|A_t| \geq \frac{\omega_{n-1}}{n}e^{\alpha_n S_p} + O(e^{-\frac{2}{n}\alpha_n t}).$$

So, by (3.4) we have

$$\begin{array}{rcl} -\frac{d|A_t|}{dt} & \geq & \alpha_n \frac{\omega_{n-1}}{n} e^{-\alpha_n t + \alpha_n S_p}(1 + O(e^{-\frac{2}{n}\alpha_n t}))(1 + O(e^{-\frac{2}{n}\alpha_n t})) \\ & = & \omega_{n-1}^{\frac{n}{n-1}} e^{-\alpha_n t + \alpha_n S_p}(1 + O(e^{-\frac{2}{n}\alpha_n t})). \end{array}$$

□

Now, we are able to construct a function sequence $\{u_\epsilon\} \subset H_0^{1,n}(M)$ which satisfies

$$\int_M F_{1,m}(u_\epsilon)dV_g > \mu(M) + \frac{\omega_{n-1}}{n}e^{\alpha_n S_p + 1 + 1/2 + \cdots + /1(n-1)}.$$

Let

$$f_\epsilon(t) = \begin{cases} C_\epsilon - \frac{(n-1)\log(1 + c_n \frac{e^{-\frac{\alpha_n}{n-1}t}}{\epsilon^{\frac{n}{n-1}}}) + \Lambda_\epsilon}{\alpha_n C_\epsilon^{\frac{1}{n-1}}} & t > t_0 = -\frac{1}{\alpha_n}\log{(L_\epsilon \epsilon)^n} \\ \frac{t}{C_\epsilon^{\frac{1}{n-1}}} & t \leq t_0 = -\frac{1}{\alpha_n}\log(L_\epsilon \epsilon)^n \end{cases} \tag{3.5}$$

where $\Lambda_\epsilon, C_\epsilon, L_\epsilon$ will be defined later (by (3.6), (3.7), (3.10)), which satisfy

i)$L_\epsilon \to +\infty$, $C_\epsilon \to +\infty$, and $L_\epsilon \epsilon \to 0$, as $\epsilon \to 0$;

ii)$C_\epsilon - \frac{(n-1)\log(1+c_n L_\epsilon^{\frac{n}{n-1}})+\Lambda_\epsilon}{\alpha_n C_\epsilon^{\frac{1}{n-1}}} = \frac{-\frac{1}{\alpha_n}\log(L_\epsilon \epsilon)^n}{C_\epsilon^{\frac{1}{n-1}}}$;

iii)$\frac{\log L_\epsilon}{C_\epsilon^{\frac{n}{n-1}}} \to 0$, as $\epsilon \to 0$.

Set $u_\epsilon = f_\epsilon(G_p)$, we have $u_\epsilon \in H^{1,n}$ because of ii).

Remark 3.1: When $M = B_1(0)$, the expression of u_ϵ will become quite simple:

$$u_\epsilon = \begin{cases} C_\epsilon - \frac{(n-1)\log(1+c_n|\frac{x}{\epsilon}|^{\frac{n}{n-1}})+\Lambda_\epsilon}{\alpha_n C_\epsilon^{\frac{1}{n-1}}} & |x| \le L\epsilon \\ -\frac{\log r^n}{\alpha_n C_\epsilon^{\frac{1}{n-1}}} & |x| > L\epsilon. \end{cases}$$

We have

$$\begin{aligned} \int_{A_{t_0}^c} |\nabla u_\epsilon|^n dV_g &= \int_{A_{t_0}^c} \frac{1}{C_\epsilon^{\frac{n}{n-1}}} |\nabla G_p|^n dV_g \\ &= \int_{\partial A_{t_0}} \frac{t_0}{C_\epsilon^{\frac{n}{n-1}}} |\nabla G_p|^{n-2} \frac{\partial G_p}{\partial n} \\ &= \frac{-\log(L_\epsilon \epsilon)^n}{\alpha_n C_\epsilon^{\frac{n}{n-1}}}. \end{aligned}$$

and

$$\begin{aligned} \int_{A_{t_0}} |\nabla u_\epsilon|^n dV_g &= \int_{t_0}^\infty |f'|^n |\nabla G_p|^n \int_{\partial A_t} \frac{1}{|\nabla G_p|} dS_t dt \\ &= \int_{t_0}^\infty |f'|^n \int_{\partial A_t} |\nabla G_p|^{n-1} dS_t dt \\ &= \int_{t_0}^\infty |f'|^n dt. \end{aligned}$$

Since

$$\begin{aligned} \int_{t_0}^\infty |f'(t)|^n dt &= \int_{t_0}^\infty \left| \frac{c_n \frac{e^{\frac{-\alpha_n t}{n-1}}}{\epsilon^{\frac{n}{n-1}}}}{C_\epsilon^{\frac{1}{n-1}} (1+c_n \frac{e^{-\frac{\alpha_n t}{n-1}}}{\epsilon^{\frac{n}{n-1}}})} \right|^n dt \\ &= \frac{n-1}{\alpha_n C_\epsilon^{\frac{n}{n-1}}} \int_0^{c_n L_\epsilon^{\frac{n}{n-1}}} \frac{u^{n-1}}{(1+u)^n} du \qquad (u = c_n \frac{e^{\frac{-\alpha_n t}{n-1}}}{\epsilon^{\frac{n}{n-1}}}) \\ &= \frac{n-1}{\alpha_n C_\epsilon^{\frac{n}{n-1}}} \int_0^{c_n L_\epsilon^{\frac{n}{n-1}}} \frac{((1+u)-1)^{n-1}}{(1+u)^n} du \\ &= \frac{n-1}{\alpha_n C_\epsilon^{\frac{n}{n-1}}} \sum_{k=0}^{n-2} \frac{C_{n-1}^k (-1)^{n-1-k}}{n-k-1} \\ &\quad + \frac{n-1}{\alpha_n C_\epsilon^{\frac{n}{n-1}}} \log(1 + c_n L^{\frac{n}{n-1}}) + O(\frac{1}{L^{\frac{n}{n-1}} C_\epsilon^{\frac{n}{n-1}}}) \\ &= -\frac{n-1}{\alpha_n C_\epsilon^{\frac{n}{n-1}}} (1 + 1/2 + 1/3 + \cdots + 1/(n-1)) \\ &\quad + \frac{n-1}{\alpha_n C_\epsilon^{\frac{n}{n-1}}} \log(1 + c_n L_\epsilon^{\frac{n}{n-1}}) + O(\frac{1}{L_\epsilon^{\frac{n}{n-1}} C_\epsilon^{\frac{n}{n-1}}}), \end{aligned}$$

where we use the fact

$$-\sum_{k=0}^{n-2} \frac{C_{n-1}^k (-1)^{n-1-k}}{n-k-1} = 1 + \frac{1}{2} + \cdots + \frac{1}{n-1}.$$

We get

$$\begin{aligned}\textstyle\int_M |\nabla u_\epsilon|^n dV_g &= \tfrac{1}{\alpha_n {C_\epsilon}^{\frac{n}{n-1}}}\{-(n-1)(1+1/2+\cdots+1/(n-1)) \\ &\quad +(n-1)\log(1+c_n {L_\epsilon}^{\frac{n}{n-1}}) - \log(L_\epsilon \epsilon)^n + O({L_\epsilon}^{-\frac{n}{n-1}})\}.\end{aligned}$$

Set $\int_M |\nabla u_\epsilon|^n dV_g = 1$, we have

$$\begin{aligned}\alpha_n {C_\epsilon}^{\frac{n}{n-1}} &= -(n-1)(1+1/2+\cdots+1/(n-1)) \\ &\quad + \log\tfrac{(1+c_n {L_\epsilon}^{\frac{n}{n-1}})^{n-1}}{{L_\epsilon}^n} - \log\epsilon^n + O({L_\epsilon}^{-\frac{n}{n-1}}) \\ &= -(n-1)(1+1/2+\cdots+1/(n-1)) + \\ &\quad \log\tfrac{\omega_{n-1}}{n} - \log\epsilon^n + O({L_\epsilon}^{-\frac{n}{n-1}}).\end{aligned} \tag{3.6}$$

By ii), we get

$$\Lambda_\epsilon = -(n-1)(1+1/2+\cdots+1/(n-1)) + O({L_\epsilon}^{-\frac{n}{n-1}}). \tag{3.7}$$

Next, we compute $\int_M e^{\alpha_n |u_\epsilon|^{\frac{n}{n-1}}} dV_g$.

Clearly, $\varphi(t) = |1-t|^{\frac{n}{n-1}} - \frac{n}{n-1}t$ is increasing when $0 \le t \le 1$ and decreasing when $t \le 0$, then

$$|1-t|^{\frac{n}{n-1}} \ge 1 - \frac{n}{n-1}t \;\; when \;\; t < 1,$$

thus when $t > t_0$, we have

$$\begin{aligned}\alpha_n f_\epsilon^{\frac{n}{n-1}}(t) &= \alpha_n {C_\epsilon}^{\frac{n}{n-1}} \Big|1 - \frac{(n-1)\log(1+c_n \frac{e^{-\frac{\alpha_n}{n-1}t}}{\epsilon^{\frac{n}{n-1}}}) + \Lambda_\epsilon}{\alpha_n {C_\epsilon}^{\frac{n}{n-1}}}\Big|^{\frac{n}{n-1}} \\ &\ge \alpha_n {C_\epsilon}^{\frac{n}{n-1}} \Big(1 - \tfrac{n}{n-1}\frac{(n-1)\log(1+c_n \frac{e^{-\frac{\alpha_n}{n-1}t}}{\epsilon^{\frac{n}{n-1}}}) + \Lambda_\epsilon}{\alpha_n {C_\epsilon}^{\frac{n}{n-1}}}\Big).\end{aligned} \tag{3.8}$$

Applying Lemma 3.1 ,we have

$$\begin{aligned}\textstyle\int_{A_{t_0}} e^{\alpha_n |u_\epsilon|^{\frac{n}{n-1}}} dV_g &= \textstyle\int_{t_0}^\infty e^{f(t)^{\frac{n}{n-1}}} \int_{\partial A_t} \frac{dS_t}{|\nabla G_p|} dt \\ &\ge \textstyle\int_{t_o}^\infty e^{\alpha_n {C_\epsilon}^{\frac{n}{n-1}} - n\log(1+c_n \frac{e^{\frac{-\alpha_n}{n-1}t}}{\epsilon^{\frac{n}{n-1}}}) - \frac{n}{n-1}\Lambda_\epsilon} \\ &\quad \times e^{-\alpha_n t + \alpha_n S_p} \omega_{n-1}^{\frac{n}{n-1}} (1 + O(e^{-\frac{2\alpha_n}{n}t})) \\ &= (1 + O(e^{-\frac{2\alpha_n}{n}t_0})) e^{\alpha_n {C_\epsilon}^{\frac{n}{n-1}} + \alpha_n S_p - \frac{n}{n-1}\Lambda_\epsilon} \\ &\quad \times \textstyle\int_{t_0}^\infty \frac{e^{-\alpha_n t}}{(1+c_n \frac{e^{\frac{-\alpha_n}{n-1}t}}{\epsilon^{\frac{n}{n-1}}})^n} \omega_{n-1}^{\frac{n}{n-1}} dt.\end{aligned}$$

Clearly,

$$\begin{aligned}
\int_{t_0}^{\infty} \frac{e^{-\alpha_n t}}{(1+c_n \frac{e^{\frac{-\alpha_n t}{n-1}}}{\epsilon^{\frac{n}{n-1}}})^n} \omega_{n-1}^{\frac{n}{n-1}} dt &= (n-1)\epsilon^n \int_0^{c_n L_\epsilon^{\frac{n}{n-1}}} \frac{u^{n-2}}{(1+u)^n du} \qquad (u = c_n \frac{e^{-\frac{\alpha_n}{n-1}t}}{\epsilon^{\frac{n}{n-1}}}) \\
&= (n-1)\epsilon^n \int_0^{c_n L_\epsilon^{\frac{n}{n-1}}} \frac{((u+1)-1)^{n-2}}{(1+u)^n} du \\
&= \epsilon^n (1 + O(L_\epsilon^{-\frac{n}{n-1}})).
\end{aligned}$$

Here, we used the fact

$$\sum_{k=0}^{m} \frac{(-1)^{m-k}}{m-k+1} C_m^k = \frac{1}{m+1}.$$

Then, applying (3.6) and (3.7), we have

$$\begin{aligned}
\int_{A_{t_0}} e^{\alpha_n |u_\epsilon|^{\frac{n}{n-1}}} dV_g &\geq \frac{\omega_{n-1}}{n} e^{\alpha_n S_p + 1 + 1/2 + \cdots + 1/(n-1)} \\
&\quad \times (1 + O((L_\epsilon \epsilon)^2))(1 + O(L_\epsilon^{-\frac{n}{n-1}})) \\
&= \frac{\omega_{n-1}}{n} e^{\alpha_n S_p + 1 + 1/2 + \cdots + 1/(n-1)} + \\
&\quad + O((L_\epsilon \epsilon)^2) + O(L_\epsilon^{-\frac{n}{n-1}}).
\end{aligned}$$

From iii), it follows that $\frac{C_\epsilon}{2} < u_\epsilon|_{A_{t_0}} < 2C_\epsilon$. Hence, we have

$$\int_{A_{t_0}} g_m(u_\epsilon) = O(C_\epsilon^{\frac{mn}{n-1}} L^2 \epsilon^2).$$

Moreover, we have

$$\int_{A_{t_0}^c} (e^{\alpha_n |u_\epsilon|^{\frac{n}{n-1}}} - g_m(u_\epsilon)) dV_g \geq \mu(M) - \mu(A_{t_0}) + \int_{A_{t_0}^c} \frac{\alpha_n^{m+1}}{(m+1)!} \left| \frac{G_p}{C_\epsilon^{\frac{1}{n-1}}} \right|^{\frac{n(m+1)}{n-1}} dV_g,$$

thus, we get

$$\begin{aligned}
&\int_M (e^{\alpha_n |u_\epsilon|^{\frac{n}{n-1}}} - g_m(u_\epsilon)) dV_g \\
&\quad \geq \mu(M) + \frac{\omega_{n-1}}{n} e^{\alpha_n S_p + 1 + 1/2 + \cdots + 1/(n-1)} + \\
&\quad \left(\int_{A_{t_0}^c} \frac{\alpha_n^{m+1}}{(m+1)!} \left| \frac{G_p}{C_\epsilon^{\frac{1}{n-1}}} \right|^{\frac{n(m+1)}{n-1}} dV_g + O(C_\epsilon^{\frac{mn}{n-1}} L^2 \epsilon^2) + O((L_\epsilon \epsilon)^2) + O(L_\epsilon^{-\frac{n}{n-1}}) \right) \qquad (3.9) \\
&\quad = \mu(M) + \frac{\omega_{n-1}}{n} e^{\alpha_n S_p + 1 + 1/2 + \cdots + 1/(n-1)} + \\
&\quad\quad C_\epsilon^{-\frac{n(m+1)}{(n-1)^2}} \left(\int_{A_{t_0}^c} \frac{|\alpha_n G_p^{\frac{n}{n-1}}|^{m+1}}{(m+1)!} dV_g + O(C_\epsilon^{\frac{n(m+1)}{(n-1)^2}} (L_\epsilon \epsilon)^2) + O(C_\epsilon^{\frac{n(m+1)}{(n-1)^2}} L_\epsilon^{-\frac{n}{n-1}}) \right).
\end{aligned}$$

Let

$$L_\epsilon = (-\log \epsilon)^{m+1}, \qquad (3.10)$$

then as $\epsilon \to 0$, we have

$$C_\epsilon^{\frac{n(m+1)}{(n-1)^2}} L_\epsilon^{-\frac{n}{n-1}} \to 0,\ C_\epsilon^{\frac{n(m+1)}{(n-1)^2}} (L_\epsilon \epsilon)^2 \to 0 \ and\ \frac{\log L}{C_\epsilon^{\frac{n}{n-1}}} \to 0.$$

Therefore, i),ii),iii) holds and we can conclude from (3.9) that

$$\int_M F_{1,m}(u_\epsilon) dV_g > \mu(M) + \frac{\omega_{n-1}}{n} e^{\alpha_n S_p + 1 + 1/2 + \cdots + 1/(n-1)}.$$

References

[1] L. Carleson and S. Y. A. Chang: on the existence of an extremal function for an inequality of J.Moser. Bull. Sc. Math., **110**:113-127, 1986.

[2] D.G. de Figueiredo, J.M. do O', B.Ruf: On a inequality by N.Trudinger and J.Moser and related elliptic equations. Comm. Pure. Appl. Math., **55**:135-152, 2002.

[3] M. Flucher: Extremal functions for Trudinger-Moser inequality in 2 dimensions. Comment. Math. Helv., **67**:471-497, 1992.

[4] S. Kichenassamy and L. Veron: Singular solutions of the p-laplace equation. Math. Ann., **275**:599-615, 1986.

[5] J. Moser: A sharp form of an Inequality by N.Trudinger. Ind. Univ. Math. J., **20**:1077-1091, 1971.

[6] Y. Li: Moser-Trudinger inequality on manifold of dimesion two. J. P. D. E. **14**:163–192, 2001.

[7] Y. Li: The existence of the extremal function of Moser-Trudinger inequality on compact Riemannian manifolds. To appear in Sci. China Ser. A.

[8] K.C. Lin: Extremal functions for Moser's inequality. Trans. Amer. Math. Sco., **348**:2663-2671, 1996.